



Published in final edited form as:

*Psychol Med.* 2011 November ; 41(11): 2297–2304. doi:10.1017/S0033291711000614.

## Schizophrenia Patients are Impaired in Empathic Accuracy

Junghee Lee<sup>1,2</sup>, Jamil Zaki<sup>3</sup>, Philippe-Olivier Harvey<sup>1,2</sup>, Kevin Ochsner<sup>3</sup>, and Michael F. Green<sup>1,2</sup>

<sup>1</sup>Semel Institute for Neuroscience and Human Behavior, UCLA, Los Angeles, CA

<sup>2</sup>VA Greater Los Angeles Healthcare System, Los Angeles, CA

<sup>3</sup>Department of Psychology, Columbia University, New York, NY

### Abstract

**Background**—Empathy is crucial for successful social relationships. Despite its importance for social interactions, little is known about empathy in schizophrenia. This study investigated the degree to which schizophrenia patients can accurately infer the affective state of another person (i.e. empathic accuracy).

**Methods**—Thirty schizophrenia patients and 22 healthy controls performed an Empathic Accuracy task on which they continuously rated the affective state of another person shown in a video (referred to as the “target”). These ratings were compared with the target’s own continuous self-rating of affective state; empathic accuracy was defined as the correlation between participants’ ratings and the targets’ self-ratings. A separate line-tracking task was administered to measure motoric / attentional factors that could account for group differences in performance. Participants’ self-rated empathy was measured using the Interpersonal Reactivity Index, and targets’ self-rated emotional expressivity was measured using the Berkeley Expressivity Questionnaire.

**Results**—Compared to controls, schizophrenia patients showed lower empathic accuracy although they performed the motoric tracking task at high accuracy. There was a significant group by target expressivity interaction such that patients showed a smaller increase in empathic accuracy with higher levels of emotional expressivity by the target, compared with controls. Patients’ empathic accuracy was uncorrelated with self-reported empathy or clinical symptoms.

**Conclusions**—Schizophrenia patients showed lower empathic accuracy than controls, and their empathic accuracy was less influenced by the emotional expressivity of the target. These findings suggest that schizophrenia patients benefit less from social cues of another person when making an empathic judgment.

### Keywords

empathic accuracy; expressivity; social cognition; schizophrenia

## INTRODUCTION

The capacity to be empathic – sharing and understanding the emotional states of others and responding appropriately to those states - is crucial for maintaining successful social

---

Correspondence: Junghee Lee, PhD. 300 Medical Plaza Room 2261, Semel Institute for Neuroscience and Human Behavior, UCLA, Los Angeles, CA 90095-6968 (jungheelee@ucla.edu).

### Financial Disclosure:

None of the authors has any conflict to disclose.

relationships (Eisenberg and Miller, 1987). As a fundamental interpersonal phenomenon, empathy comes into play in virtually all social interactions, and as such, difficulties in experiencing empathy may lead to social dysfunctions, including those that characterize severe mental illnesses such as schizophrenia and autism (Henry et al., 2008, Blair, 2005). Despite its potential importance in interpersonal interactions, little is known about empathy in schizophrenia.

Empathy is generally regarded as an ability to understand emotions and feelings of another person; it is a complex construct comprised of multiple abilities whose inter-relationships have been the subject of debate (Davis, 1983, Gallese et al., 2004, de Vignemont and Singer, 2006, Singer, 2006, Singer and Lamm, 2009, Marangoni et al., 1995, Ickes et al., 1990, Decety and Jackson, 2004, Preston and de Waal, 2002). Two of the most studied components of empathy are the ability to share or mimic the internal affective or intentional states of others (e.g. sharing the sadness of a grieving friend) and the ability to make explicit social cognitive attributions about those mental states (“I think he’s sad.”). A third area concerns self-reported trait empathy that involves self-assessment of one’s empathic abilities (e.g. endorsing questionnaire items like, “I take on the sadness of others”). Few studies have examined empathy in schizophrenia and the initial findings suggest that patients show differences from controls in these abilities. For example, schizophrenia patients tend to mimic less than controls when presented with another person yawning or laughing (Haker and Rossler, 2009), show atypical neural activation when making explicit attributions about other’s emotions (Benedetti et al., 2009, Lee et al., 2010), and exhibit lower scores on questionnaires assessing self-reported trait empathy (Derntl et al., 2009, Benedetti et al., 2009, Montag et al., 2007, Shamay-Tsoory et al., 2007, Sparks et al., 2010).

While these studies, using diverse methods, suggest impaired empathy in schizophrenia, they provide little insight into the nature of empathic processes in schizophrenia in real-world social situations. Interactions with others in real-life involve multi-modal social cues that are typically dynamic and rapidly changing (Zaki and Ochsner, 2009). Static, unimodal stimuli (e.g. pictures of posed facial expressions of emotion) of the sort that have been employed in previous studies on empathy in schizophrenia do not fully capture the experience of reading empathic cues in a natural setting. More importantly, empathic behavior in everyday life involves not just a perceiver (i.e. a person who is empathizing) but a target as well (i.e. the person whose affective state is being shared and/or inferred). Hence, it is necessary to consider both the cues sent by the target and the ability to read these cues by the perceiver when examining the empathic behavior of schizophrenia patients (Zaki et al., 2008).

The ability to empathize with another person from naturalistic stimuli has been studied in the context of empathic accuracy, which refers to the ability to accurately judge the amount and kind of emotion experienced by another person (Levenson and Ruef, 1992, Marangoni et al., 1995, Ickes et al., 1990, Zaki et al., 2008). Empathic accuracy is important for everyday life in that inaccurate empathic judgments would lead to social misperceptions, inappropriate responses, and problems at work or school – all of which are common in schizophrenia. A typical empathic accuracy task asks participants or perceivers to continuously judge the emotional experiences of “target” individuals describing emotionally charged autobiographical events on a video clip. Empathic accuracy can be assessed by the extent to which the perceiver’s rating of the target’s emotion matches the target’s own self-rated emotional response moment to moment. Importantly, empathic accuracy depends on characteristics of both perceivers and targets. Empathic accuracy is generally high when perceivers judge the emotional experiences of targets who describe themselves as being emotional expressive (Zaki et al., 2008, Flury et al., 2009, Snodgrass et al., 1998), presumably because their emotions are more easily readable. Further, target expressivity

appears to moderate the relationship between self-reported trait empathy and empathic accuracy: high self-reported trait empathy of a perceiver predicts high empathic accuracy, but only when a perceiver was judging the affective state of highly expressive targets (Zaki et al., 2008).

In the current study, we used an empathic accuracy task to study the ability of schizophrenia patients and healthy controls to correctly assess the emotions of others using naturalistic social stimuli (Zaki et al., 2008, Zaki et al., 2009b, Zaki and Ochsner, 2009). We had three primary goals. First, we examined whether schizophrenia patients showed lower empathic accuracy compared to controls. Second, we examined whether the level of emotional expression of a target moderated group differences between patients and controls. Third, we examined whether empathic accuracy was associated with self-reported trait empathy or clinical characteristics in the schizophrenia sample.

## METHODS

### Participants

Thirty patients with schizophrenia and 22 healthy controls participated in this study. All participants received the Structured Clinical Interview for DSM-IV (SCID) Axis I Disorders (First et al., 1997) to confirm their eligibility. Schizophrenia patients were recruited from outpatient clinics at the Veterans Affairs (VA) Greater Los Angeles Healthcare System and University of California, Los Angeles and from local board and care facilities. Healthy control participants were recruited through flyers posted in the local community and website postings. Exclusion criteria for patients included: 1) substance abuse or dependence in the last six months based on the SCID (First et al., 1997), 2) current major depressive episode, 3) mental retardation based on review of medical records, 4) history of loss of consciousness for more than one hour due to head trauma, 5) an identifiable neurological disorder, or 6) insufficient fluency in English to understand the procedures based on clinician's judgment. Controls were excluded if they had: 1) history of schizophrenia or other psychotic disorder, bipolar disorder, recurrent depression, substance dependence, or any substance abuse in the last 6 months based on the SCID (First et al., 1997), 2) current major depressive episode, 3) any of the following Axis II disorders: avoidant, paranoid, schizoid, or schizotypal, based on the SCID for Axis II disorders (First et al., 1996), 4) schizophrenia or other psychotic disorder in a first-degree relative, 5) any significant neurological disorder or head injury, or 6) insufficient fluency in based on clinician's judgment.

Schizophrenia patients and healthy controls were comparable in terms of age and parental education, but not personal education (see Table 1 for demographic information). All of the patients were taking antipsychotic medications at the time of testing. All participants had normal or corrected to normal vision of at least 20/30. Using the expanded 24-item version of the Brief Psychiatric Rating Scale (BPRS; Ventura et al., 1993), clinical symptoms for patients were divided into 3 factors: thinking disturbance factor consisting of unusual thought content, hallucination, and conceptual disorganization; withdrawal-retardation factor consisting of blunted affect, emotional withdrawal, and motor retardation; and anxiety / depression factor composed of somatic concern, anxiety, depression, and guilt.

All interviewers were trained through the Treatment Unit of the Department of Veterans Affairs VISN 22 Mental Illness Research, Education, and Clinical Center (MIRECC). SCID interviewers were trained to a minimum kappa of 0.75 for key psychotic and mood items, and symptom raters were trained to a minimum intraclass correlation (ICC) of 0.80. All participants were evaluated for the capacity to give informed consent and provided written informed consent after all procedures were fully explained, according to procedures

approved by the Institutional Review Board at the VA Greater Los Angeles Healthcare System.

### Empathic Accuracy Task

The Empathic Accuracy Task was adapted from Zaki et al. (Zaki et al., 2008, Zaki et al., 2009b). It consisted of 12 video clips (6 positive valence and 6 negative valence), each lasting for 1 – 2.3 minutes (mean, 102 seconds; range 62 – 137 seconds). A detailed explanation of the development of these videos is provided elsewhere (Zaki et al., 2008, Zaki et al., 2009b). Briefly, the head and shoulders of an individual (referred to as “target”) were videotaped while he / she discussed a positive or negative autobiographical event. Immediately after the videos were filmed, targets: 1) provided continuous ratings of their own emotional experience while watching their own videos and 2) completed the 10-item Berkeley Expressivity Questionnaire (BEQ; Gross, 2000), which assess tendencies to experience and express strong emotions in general. Six positive and six negative videos of this study had equal number of male and female targets and were equated for the target’s self-reported expressivity rated by the BEQ.

For the current study, a central fixation was presented at the start of each trial. Immediately after the fixation disappeared, a video was presented in the center of a black screen. Above the video, an instruction was presented that oriented participants to the judgment they were to make (i.e. how good or bad is this person feeling?). Below the video, a 9-point rating scale was presented (1: very negative, 5: neutral, and 9: very positive). Participants were asked to continuously rate how positive or negative they believed the target was feeling at each moment using the left or right arrow keys. Each video started with the number 5 selected and participants pressed the left or right arrow key to move the number upward (toward positive) or downward (toward negative). The selected number on the scale was always highlighted so that participants could monitor their ratings of the target’s emotion.

If patients show poorer empathic accuracy than controls, it is important to verify that it is not due to the motoric rather than the empathic demands of the task. We considered this confound in two ways. First, we recorded the number of times subjects pressed the arrow keys to see if patients were making rating responses significantly less often than controls. Second, most of the participants (26 patients and 14 controls) received an additional motor tracking task that approximated the motoric demands of the empathic accuracy task but did not involve a social component. This task was designed to examine whether patients were able to sustain their attention and track a range of changes for about 2 minutes. This task included two videos (each lasting 2 minutes) that showed a thin vertical red line moving left or right at varying speeds. Similar to the empathic accuracy task, participants continuously rated the location of a moving visual target by pressing the arrow keys right or left along a 9-point scale.

### The Interpersonal Reactivity Index (IRI)

All participants completed the IRI (Davis, 1983) to assess self-reported trait empathy. The IRI is a reliable and valid way of measuring the one’s belief in one’s own empathic tendencies (Davis, 1994). The IRI consists of four subscales, each with 7 items: Fantasy (FS), Perspective Taking (PT), Empathic Concern (EC), and Personal Distress (PD). Participants responded using a 5-point Likert scale, with 1 being *does not describe me well* and 5 being *describes me very well*. The Fantasy subscale measures the tendency to imagine oneself in a fictional situation (e.g. “After seeing a play or movie, I have felt as though I were one of the characters.”). The Perspective Taking subscale assesses the tendency to adopt the point of view of others and reason about their mental state (e.g. “I try to look at everybody’s side of a disagreement before I make a decision.”). The Empathic Concern

subscale measures the tendency to experience emotions in response to others and/or sympathy and concern for them (e.g. “I am often quite touched by things that I see happen.”). The Personal Distress subscale assesses the tendency to experience distress or discomfort in response to others’ misfortune (e.g. “In emergency situations, I feel apprehensive and ill-at-ease.”).

### Statistical Analyses

Data reduction and time series analysis for the empathic accuracy and motoric tracking tasks were conducted using Matlab (Mathworks). Continuous affect ratings were converted into a time series of sequential values – one number for every two seconds of video. Specifically, the average rating was determined for each 2-second epoch for each participant, and these values served as data points in subsequent time series analyses. To calculate empathic accuracy, participants’ continuous ratings across these 2-second epochs were correlated with the target’s own continuous ratings across the same epochs for each video. The resulting correlation coefficient ( $r$ ) between two time series is the measure of empathic accuracy. For the motoric tracking task, the participants’ responses to the movement of the line were correlated with the actual line movement for each video. Before conducting any statistical analyses, the individual correlation coefficients for both tasks were converted into  $z$  scores, which were used in subsequent analyses. To compare group difference on the empathic accuracy task,  $z$ -scores were summed for positive and negative valence separately and a  $2 \times 2$  repeated measures ANOVA was performed with valence as a within-subject factor and group as a between-subject factor. For the motoric tracking task, a  $t$ -test for independent samples was used to examine group differences. To examine the effect of the target’s expressivity on empathic accuracy score, the mixed linear model was performed with the expressivity and group as fixed effects and subject as a random effect. To examine whether impaired empathic accuracy of patients is related to other characteristics, we examined its correlation with the four subscales of the IRI and clinical symptoms measured by BPRS.

## RESULTS

Figure 1 shows the empathic accuracy scores ( $r$ ) for each group. The patients’ empathic accuracy scores for both positive and negative valence were significantly above zero ( $t_{29} = 9.48, p < .001$  and  $t_{29} = 7.66, p < .001$ , respectively). For the  $2 \times 2$  ANOVA, the main effects of group ( $F_{1,50} = 12.88, p < .01$ ) and valence ( $F_{1,50} = 7.15, p < .01$ ) were both significant. The interaction was not significant. Overall, patients showed lower accuracy for rating the target’s emotion and both groups showed better accuracy for positive valence. We then examined possible confounds for this significant group difference in empathic accuracy by assessing whether groups differed in the number of manual responses during the task. There was a significant main effect of valence ( $F_{1,50} = 14.27, p < .001$ ); both groups responded more to negative than positive videos. However, neither the group nor the interaction was significant (patients: positive =  $13.3 \pm 17.6$ , negative =  $21.8 \pm 29.9$ ; controls, positive =  $7.4 \pm 5.5$ , negative =  $10.3 \pm 6.9$ ) although patients responded more frequently than controls. We also examined whether patients accurately performed the motoric tracking task. Both groups tracked the line with extremely high accuracy ( $r = .90, SD = .16$ ;  $r = .97, SD = .02$ , for patients and controls, respectively). This difference was significant ( $t_{36} = -3.63, p < .01$ ), but the high level of accuracy in patients indicates that the motoric aspects of the empathic accuracy task are unlikely to account for their lower empathic accuracy.

Next we considered the effect of target expressivity on empathic accuracy using a mixed linear model. We found a significant main effect of expressivity ( $F_{1,566} = 21.78, p < .001$ ) and a significant group by expressivity interaction ( $F_{1,566} = 3.18, p < .05$ ) indicating that the expressivity of the target was associated with higher empathic accuracy scores in both groups but this relationship was significantly smaller in schizophrenia patients. Figure 2



shows the mean of standardized empathic accuracy scores from patient group and control group for each video as a function of the target's expressivity.

Finally, we examined whether the empathic accuracy of schizophrenia patients was related to: 1) their self-reported empathy and 2) clinical symptoms. The mean (SD) trait empathy subscale scores and the three clinical symptom factors are shown in Table 1. No significant relationship was observed between empathic accuracy score and the four IRI subscales. Similarly, empathic accuracy was not correlated with the three factors of the BPRS.

## DISCUSSION

In this study we examined one key aspect of empathy in schizophrenia; namely, empathic accuracy, which refers to the ability to make accurate empathic judgments of others' emotions. We used an empathic accuracy task that required participants to track, in real-time, the fluctuating contours of a target's emotions in a manner that approximates the need to track emotion in real world social interactions. Three key findings were obtained. First, compared with controls, schizophrenia patients showed reduced empathic accuracy across both positive and negative video clips, indicating that schizophrenia patients are less accurate at inferring the affective state of another person. The impaired performance of patients is less likely attributable to impairments in sustained attention or motor abilities, given that schizophrenia patients tracked a dynamically moving non-social visual stimulus over 2 minutes with high accuracy. Second, although both groups showed greater empathic accuracy for highly expressive targets, this effect was significantly smaller in schizophrenia patients. Third, empathic accuracy was not related to their self-reported belief in their empathic ability or clinical symptoms within the schizophrenia group. Taken together, these results suggest that schizophrenia patients are likely to have reduced empathic ability compared with controls interacting with others in everyday life, and these differences are greatest when interacting with highly emotionally expressive people.

Despite the rapid emergence of research on social cognition in schizophrenia, there is a relative paucity of appropriate measures for studying empathic processes in this population (Green et al., 2008). Empathy is a multi-faceted construct and the few previous studies on empathy in schizophrenia focused on certain aspects, such as low trait empathy or deficits in mimicking the affect of another person (Haker and Rossler, 2009, Derntl et al., 2009, Benedetti et al., 2009, Montag et al., 2007, Shamay-Tsoory et al., 2007, Sparks et al., 2010). However, the ability to accurately understand the affective state of another person is an important component of empathy that had not been previously studied in schizophrenia. The current study showed that schizophrenia patients were impaired in empathic accuracy; that is, they were less accurate at tracking the positive and negative affective state of another person compared to controls.

The empathic accuracy task used in this study involved the ability to monitor the valence of others' affect, as opposed to the ability to decode the specific content of targets' thoughts and feelings; both of these approaches provide important and complementary information about empathic abilities. Whereas content-driven approach accuracy tasks can provide more specific insights about what types of thoughts and feelings perceivers accurately understand, they tend to be highly dependent on verbal ability (Davis and Kraus, 1997), and often require somehow subjective coding of accuracy based on targets' and perceivers' verbal reports. By contrast, a valence-driven approach provides more 'coarse' information about accuracy for positive and negative affect, but also produces a highly tractable, quantitative operationalization of accuracy that does not depend on target's or perceivers' specific verbal content (Zaki et al., 2009a). Further, both valence- and content-focused measures of empathic accuracy are associated with functionally important outcomes, such as relationship

quality and social support (Zaki and Ochsner, in press, Bartz et al., in press, Carton et al., 1999). As such, impairments in either types of accuracy are likely to provide meaningful sources of information about poor social functioning in schizophrenia.

In this study empathic accuracy in both schizophrenia patients and controls were not associated with self-reported trait empathy. The lack of association between empathic accuracy and trait empathy may seem surprising and counterintuitive. However, such dissociation between empathic accuracy behavior and self-report empathy is quite common (Zaki et al., 2008, Levenson and Ruef, 1992, Ickes et al., 2000). Self-reported trait empathy concerns the belief about one's own empathic characteristics, whereas empathic accuracy measures how well a person understands the affective state of another person. Further, individuals with high-self-reported trait empathy are often no better at understanding the affective state of another person than people with low self-rated empathy (Hall, 1979, Ickes et al., 1990, Levenson and Ruef, 1992, Ames and Kammrath, 2004). Thus, self-reported trait empathy appears to reflect one's own belief in his / her empathic characteristic, as opposed to the strength or accuracy of his / her empathic ability. The current findings suggest that the empathic accuracy diverges from the self-reported trait empathy in schizophrenia as well as in healthy individuals and further empathizes that empathy is indeed a multi-faceted psychological construct. Further studies will help us better understand the relationship among diverse aspects of empathy in schizophrenia.

The current study advances prior work on empathy in schizophrenia in two ways. First, this study employed an experimental method that assesses the naturalistic processes of empathic judgment. In real life we often encounter dynamically fluctuating emotional experiences of another person and make moment-to-moment judgments on the transient affective state over time. It will be possible to employ this method in studies with other psychotic disorders that may also have difficulties with empathic processes to examine whether impaired empathic accuracy is specific to schizophrenia. Second, the empathic accuracy paradigm provides a clear performance metric of empathic accuracy that can be used for relating task performance to other individual difference variables such as other aspects of empathy, theory of mind, and non-social cognition.

What would contribute to lower empathic accuracy in schizophrenia? People generally have higher empathic accuracy when they interact with another person who is highly expressive (Zaki et al., 2009a, Zaki et al., 2008). While both groups were more accurate with highly expressive individuals, schizophrenia patients were significantly less able to benefit from the expressivity of the target. The reduced ability of schizophrenia patients to benefit from an expressive target could be partly due to impaired early perceptual abnormalities in schizophrenia. Previous studies with healthy individuals demonstrated that people rely more on verbal social cues than visual cues in reaching empathic accuracy judgments (Zaki et al., 2009a, Gesn and Ickes, 1999). Perhaps schizophrenia patients had difficulty detecting subtle changes of auditory information from an expressive target, resulting in lower empathic accuracy. Along these lines, schizophrenia patients show impaired early auditory processing with tone matching and this impairment has been associated with impaired prosody detection (Leitman et al., 2005, Leitman et al., 2007). Future research could determine whether lower empathic accuracy in schizophrenia is associated with early perceptual abnormalities and whether schizophrenic patients have more trouble inferring emotion from verbal versus non-verbal cues.

This study is the first, to our knowledge, to assess empathic accuracy in schizophrenia. The current findings provide a foundation for further exploration. Low empathic accuracy of schizophrenia patients found in this study may be used to identify neural correlates of empathic understanding in schizophrenia patients. Recent studies with healthy individuals

showed that higher empathic accuracy is related to increased activations in several brain regions including the medial prefrontal cortex (Zaki et al., 2009b, Zaki et al., 2010). Finally, the current study has some limitations. All of the patients were taking antipsychotic medications. With first-episode patients without medication history, it will be possible to determine the potential effect of pharmacological treatment on empathic accuracy in schizophrenia. In addition, this study did not include non-social neurocognitive measures or assess community functioning. With a larger sample and broader range of assessments, it will be possible to determine the relationship between reduced empathic accuracy and non-social cognition, or whether empathic accuracy is related to certain domains of community functioning, such as social connectedness or vocational success.

## Acknowledgments

Support for this study came from NIMH Grant MH 043292 (MFG), NIDA grant DA022541 (KO), NIMH Grant MH076137 (KO), and Autism Speaks Grant 4787 (JZ). The authors wish to thank Mark R. McGee, Shelly Crosby, and Cory Tripp for assistance in data collection.


## References

- Ames DR, Kammrath LK. Mind-reading and metacognition: Narcissism, not actual competence, predicts self-estimated ability. *Journal of Nonverbal Behavior*. 2004; 24:187–209.
- Bartz JA, Zaki J, Bolger N, Hollander E, Ludwig N, Kolevzon A, Ochsner K. Oxytocin selectively improves empathic accuracy. *Psychological Science*. in press.
- Benedetti F, Bernasconi A, Bosia M, Cavallaro R, Dallaspesza S, Falini A, Poletti S, Radaelli D, Riccaboni R, Scotti G, Smeraldi E. Functional and structural brain correlates of theory of mind and empathy deficits in schizophrenia. *Schizophrenia Research*. 2009; 114:154–60. [PubMed: 19632816]
- Blair RJR. Responding to the emotions of others: Dissociating forms of empathy through the study of typical and psychiatric population. *Consciousness and Cognition*. 2005; 14:698–718. [PubMed: 16157488]
- Carton J, Kessler E, Pape C. Nonverbal decoding skills and relationship to well-being in adults. *Journal of Nonverbal Behavior*. 1999; 23:91–100.
- Davis M. Measuring individual differences in empathy: Evidence for multidimensional approach. *Journal of Personality and Social Psychology*. 1983; 44:113–126.
- Davis, MH. *Empathy: A social psychological approach*. Colorado: Westview Press; 1994.
- Davis, MH.; Kraus, LA. *Personality and empathic accuracy*. In: Ickes, W., editor. *Empathic Accuracy*. New York: Guilford; 1997.
- de Vignemont F, Singer T. The empathic brain: how, when and why? *Trends in Cognitive Science*. 2006; 10:435–41.
- Decety J, Jackson PL. The functional architecture of human empathy. *Behavioral and Cognitive Neuroscience Review*. 2004; 3:71–100.
- Derntl B, Finkelmeyer A, Toygar TK, Hulsmann A, Schneider F, Falkenberg DI, Habel U. Generalized deficit in all core components of empathy in schizophrenia. *Schizophrenia Research*. 2009; 108:197–206.
- Eisenberg N, Miller PA. The relation of empathy to prosocial and related behaviors. *Psychological Bulletin*. 1987; 101:91–119. [PubMed: 3562705]
- First, MB.; Gibbon, M.; Spitzer, RL.; Williams, JBW.; Benjamin, L. *Structured Clinical Interview for DSM-IV Axis I Personality Disorders*. New York: State Psychiatric Institute; 1996.
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. *Structured Clinical Interview for DSM-IV Axis I Disorders - Patient Edition*. New York: State Psychiatric Institute; 1997.
- Flury JM, Ickes W, Schweinle W. The borderline empathy effect: do high BPD individuals have greater empathic ability? Or are they just more difficult to “read”? *Journal of Research in Personality*. 2009; 42:312–332.



- Gallese V, Keysers C, Rizzolatti G. A unifying view of the basis of social cognition. *Trends in Cognitive Science*. 2004; 8:396–403.
- Gesn PR, Ickes W. The development of meaning contexts for empathic accuracy: channels and sequence effects. *Journal of Personality and Social Psychology*. 1999; 77:746–761.
- Green MF, Penn DL, Bentall R, Carpenter WT, Gaebel W, Gur RC, Kring AM, Park S, Silverstein SM, Heinssen R. Social cognition in schizophrenia: an NIMH workshop on definitions, assessment, and research opportunities. *Schizophr Bull*. 2008; 34:1211–1220. [PubMed: 18184635]
- Gross, J. The Berkeley Expressivity Questionnaire. In: Maltby, J.; Lewis, CA.; Hill, A., editors. *Commissioned reviews on 300 psychological tests*. Lampeter, Wales: Edwin Mellen Press; 2000.
- Haker H, Rossler W. Empathy in schizophrenia: impaired resonance. *European Archives of Psychiatry and Clinical Neuroscience*. 2009; 259:352–61. [PubMed: 19377866]
- Hall, JA. Gender, gender roles, and nonverbal communication skills. In: Rosenthal, R., editor. *Skills in Nonverbal Communication*. Cambridge, MA: Oelgeschlager, Gunn, & Hain; 1979.
- Henry JD, Bailey PE, Rendell PG. Empathy, social functioning and schizotypy. *Psychiatry Res*. 2008; 160:15–22. [PubMed: 18511132]
- Ickes W, Buysse AHP, Rivers K, Erickson J, Hancock M, Kelleher J, Gesn PR. On the difficulty of distinguishing “good” and “poor” perceivers: A social relations analysis of empathic accuracy data. *Personal Relationships*. 2000; 7:219–234.
- Ickes W, Stinson L, Bissonnette V, Garcia S. Naturalistic social cognition: Empathic accuracy in mixed-sex dyads. *Journal of Personality & Social Psychology*. 1990; 59:730–742.
- Lee SJ, Kang DH, Kim CW, Gu BM, Park JY, Choi CH, Shin NY, Lee JM, Kwon JS. Multi-level comparison of empathy in schizophrenia: An fMRI study of a cartoon task. *Psychiatry Research: Neuroimaging*. 2010; 181:121–129.
- Leitman DI, Foxe JJ, Butler PD, Saperstein A, Revheim N, Javitt DC. Sensory contributions to impaired prosodic processing in schizophrenia. *Biological Psychiatry*. 2005; 58:56–61. [PubMed: 15992523]
- Leitman DI, Hoptman MJ, Foxe JJ, Saccente E, Wylie GR, Nierenberg J, Jalbrzikowski M, Lim KO, Javitt DC. The neural substrates of impaired prosodic detection in schizophrenia and its sensorial antecedents. *American Journal of Psychiatry*. 2007; 164:474–82. [PubMed: 17329473]
- Levenson RW, Ruef AM. Empathy: A physiological substrate. *Journal of Personality and Social Psychology*. 1992; 63:234–246. [PubMed: 1403614]
- Marangoni C, Garcia S, Ickes W, Teng G. Empathic accuracy in a clinically relevant setting. *Journal of Personality and Social Psychology*. 1995; 68:854–69.
- Montag C, Heinz A, Kunz D, Gallinat J. Self-reported empathic abilities in schizophrenia. *Schizophrenia Research*. 2007; 92:85–9. [PubMed: 17350225]
- Preston SD, de Waal FB. Empathy: Its ultimate and proximate bases. *Behavioral and Brain Sciences*. 2002; 25:1–20. discussion 20–71. [PubMed: 12625087]
- Shamay-Tsoory SG, Shur S, Harari H, Levkovitz Y. Neurocognitive basis of impaired empathy in schizophrenia. *Neuropsychology*. 2007; 21:431–8. [PubMed: 17605576]
- Singer T. The neuronal basis and ontogeny of empathy and mind reading: review of literature and implications for future research. *Neuroscience and Biobehavioral Reviews*. 2006; 30:855–63.
- Singer T, Lamm C. The social neuroscience of empathy. *Annals of the New York Academy of Sciences*. 2009; 1156:81–96. [PubMed: 19338504]
- Snodgrass SE, Hecht MA, Ploutz-Snyder R. Interpersonal sensitivity: expressivity or perceptivity? *Journal of Personality and Social Psychology*. 1998; 74:238–49. [PubMed: 9457785]
- Sparks A, McDonald S, Lino B, O'Donnell M, Green MJ. Social cognition, empathy and functional outcome in schizophrenia. *Schizophrenia Research*. 2010; 122:172–8. [PubMed: 20609567]
- Ventura J, Lukoff D, Nuechterlein KH, Liberman RP, Green MF, Shaner A. Brief Psychiatric Rating Scale (BPRS) expanded version: scales, anchor points, and administration manual. *International Journal of Methods in Psychiatric Research*. 1993; 3:227–243.
- Zaki J, Bolger N, Ochsner K. It takes two: the interpersonal nature of empathic accuracy. *Psychological Science*. 2008; 19:399–404. [PubMed: 18399894]


- Zaki J, Bolger N, Ochsner K. Unpacking the informational bases of empathic accuracy. *Emotion*. 2009a; 9:478–87. [PubMed: 19653768]
- Zaki J, Hennigan K, Weber J, Ochsner KN. Social cognitive conflict resolution: contributions of domain-general and domain-specific neural systems. *Journal of Neuroscience*. 2010; 30:8481–8. [PubMed: 20573895]
- Zaki J, Ochsner K. The need for a cognitive neuroscience of naturalistic social cognition. *Annals of the New York Academy of Sciences*. 2009; 1167:16–30. [PubMed: 19580548]
- Zaki J, Ochsner K. Re-integrating the study of accuracy into social cognition research. *Psychological Inquiry*. in press.
- Zaki J, Weber J, Bolger N, Ochsner K. The neural bases of empathic accuracy. *Proceedings of the National Academy of Sciences, U S A*. 2009b; 106:11382–11387.



**Figure 1.**

Empathic accuracy score ( $r$ ) of schizophrenia patients and controls.

The mean empathic accuracy scores ( $r$ ) of patients and controls are shown for positive and negative video clips. Schizophrenia patients showed overall lower empathic accuracy scores. Both groups showed better accuracy for understanding positive emotional experiences of a target than negative emotional experiences. Values are presented as mean (standard error).



**Figure 2.** Empathic accuracy of patient group and control group as a function of a target expressivity. The mean of standardized empathic accuracy scores from patient group and control group for each video is plotted as a function of the target’s expressivity. Both groups showed better accuracy for understanding emotional experiences of a highly expressive target; but this effect is much smaller in patients.

**Table 1**

Demographics of schizophrenia patients and healthy controls

	Schizophrenia patients	Healthy controls	Statistics
Age	46.1 (12.1)	44.3 (8.7)	$t_{50} = .59$ , NS
Education (yrs)	12.8 (1.3)	14.7 (1.7)	$t_{50} = -4.47$ , $p < .001$
Parental education (yrs)	11.6 (2.8)	13.0 (2.7)	$t_{48} = -1.83$ , NS
Gender (female / male)	5 / 25	5 / 17	$\chi^2 = .30$ , NS
BPRS (factor totals)			
Thinking Disturbance	5.9 (3.0)	NA	
Withdrawal / Retardation	5.3 (2.3)	NA	
Anxiety / Depression	8.4 (2.9)	NA	
Interpersonal Reactivity Index			
Fantasy	12.1 (4.4)	12.9 (6.1)	$t_{50} = -.48$ , NS
Perspective Taking	14.3 (4.1)	16.7 (4.2)	$t_{50} = -2.02$ , $p < .05$
Empathic concerns	17.1 (5.1)	20.5 (3.9)	$t_{50} = -2.55$ , $p < .05$
Personal distress	11.9 (6.2)	5.5 (3.3)	$t_{50} = 4.33$ , $p < .001$

<sup>†</sup> Values are given as mean (standard deviation).

<sup>††</sup> BPRS, Brief Psychiatric Rating Scale; NS, non significant; NA, not applicable